

### Triple Pascal Sequence Spaces

N. Subramanian<sup>1</sup> and A. Esi<sup>2</sup>

<sup>1</sup>Department of Mathematics, SASTRA University, Thanjavur-613 401, India

E-mail: nsmaths@gmail.com

<sup>2</sup>Department of Mathematics, Adiyaman University, 02040, Adiyaman, Turkey

E-mail: aesi23@hotmail.com

(Received 27 May, 2019)

In this paper, the concept of pascal triple sequence spaces are introduced and then basic topological properties of pascal triple sequence spaces are investigated.

Keywords: Pascal; triple sequence; Pascal sequence space.

#### I. INTRODUCTION

The triple pascal matrix is an infinite matrix containing the binomial coefficients as its elements. There are three ways to achieve this as either an upper-triangular matrix, a lower-triangular matrix or a symmetric matrix. The  $4 \times 4$  truncation of these are show below.

The triple upper triangular

$$U_4 = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 27 & 96 \\ 0 & 0 & 1 & 500 \\ 0 & 0 & 0 & 1 \end{pmatrix};$$

Triple lower triangular

$$L_4 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 27 & 1 & 0 \\ 1 & 96 & 500 & 1 \end{pmatrix};$$

Symmetric

$$A_4 = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 27 & 500 & 8575 \\ 1 & 96 & 3375 & 87808 \\ 1 & 250 & 15435 & 592704 \end{pmatrix}$$

These matrices have the pleasing relationship  $A_n = L_n U_n$ . It is easily seen that all three matrices have determinant 1. The elements of the symmetric triple pascal matrix are the binomial coefficients.

(i.e)  $A_{ijk} = \binom{r}{m} \binom{s}{n} \binom{t}{k} = \frac{r!}{m!(r-m)!} \frac{s!}{n!(s-n)!} \frac{t!}{k!(t-k)!}$ ,

where  $r, s, t = i + j + k$  and  $m = i, n = j, k = t$ .

In other words

$$A_{ijk} = {}_{i+j+k}C_{ijk} = \frac{(i+j+k)!}{i!j!k!}.$$

Thus the trace of  $A_n$  is given by

$$tr(A_n) = \sum_{m=0}^{r-1} \sum_{n=0}^{s-1} \sum_{k=0}^{t-1} \frac{(2m)!}{(m!)^2} \frac{(2n)!}{(n!)^2} \frac{(2k)!}{(k!)^2}$$

with the first few terms given by the sequence 1, 27, 729, 24389, ...

Let  $A_n$  be  $n \times n \times n$  matrix whose skew diagonals are successively the rows (truncated where necessary) of pascals triangle. In general,  $A_n = (a_{ijk})$ , where

$$a_{ijk} = \binom{i+j+k}{i} \binom{i+j+k}{j} \binom{i+j+k}{k} \text{ for } i, j, k = 0, 1, 2, \dots, n-1.$$

An possesses the factorization

$$A_n = L_n L_n^T \quad (1)$$

where  $L_n^T$  denotes the transpose of  $L_n$ . For the  $[ijk]^{th}$  section of element of this product is

$$\begin{aligned} &= \text{coefficient of } x^{ijk} \text{ in } (1+x)^i (1+x)^j (1+x)^k \\ &= a_{ijk} = \binom{i+j+k}{i} \binom{i+j+k}{j} \binom{i+j+k}{k}. \end{aligned}$$

Clearly

$$|L_n| = 1 \quad (2)$$

so that

$$|A_n| = |L_n L_n^T| = |L_n|^2 = 1$$

we observe that  $L_n^{-1}$  is simply related to  $L_n$ .

For example

$$L_4^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 1 & -27 & 1 & 0 \\ 1 & 96 & -500 & 1 \end{pmatrix};$$

and in general

$$L_n^{-1} = (-1)^{i+j-2k} I_{ijk} \quad (3)$$

In addition, 1 is an eigen value of  $A_n$  when  $n$  is odd and that if  $\lambda$  is an eigen value of  $A_n$  then so is  $\lambda^{-1}$ . These conjectures

$$P = [P_{mnk}^{rst}] = \begin{cases} \begin{pmatrix} r \\ m \end{pmatrix} \begin{pmatrix} s \\ n \end{pmatrix} \begin{pmatrix} t \\ k \end{pmatrix} \\ 0, \end{cases}$$

are readily verified for small values of  $n$ . In general

Let

$$P_n(\lambda) = |\lambda I_n - A_n|$$

where  $I_n$  is the  $n \times n \times n$  identity matrix. Then by (1.1),(1.2)

and (1.3)

$$\begin{aligned} P_n(\lambda) &= |\lambda L_n L_n^{-1} - L_n L_n^T| \\ &= |L_n| |\lambda L_n^{-1} - L_n^T| \\ &= \left| \left( (-1)^{i+j-2k} \lambda I_{ijk} - I_{kji} \right) \right| \\ &= (-\lambda)^n \left| \left( \lambda_{kji}^{-1} I - (-1)^{i+j-2k} I \right) \right|. \end{aligned}$$

Multiplying odd numbered rows and columns of the matrix by -1 and transposing, we get

$$P_n(\lambda) = (-\lambda)^n \left| \left( (-1)^{i+j-2k} \lambda_{ijk}^{-1} I - I_{kji} \right) \right|$$

$$P_n(\lambda) = (-\lambda)^n P_n \left( \frac{1}{\lambda} \right) \quad (4)$$

But eigen values of  $A_n$  are the roots of  $P_n(\lambda) = 0$  and thus it follows from (1.4) that if  $\lambda$  is an eigen value of  $A_n$  then so is  $\lambda^{-1}$ .

A triple sequence (real or complex) can be defined as a function  $X : \mathbb{N}^3 \rightarrow \mathbb{R}(\mathbb{C})$ , where  $\mathbb{N}, \mathbb{R}$  and  $\mathbb{C}$  denote the set of natural numbers, real numbers and complex numbers respectively. The different types of notions of triple sequence was introduced and investigated at the initial by *Bipan Hazarika et al. [1], Sahiner et al. [12,13], Esi et al. [2-9], Dutta et al. [10], Subramanian et al. [14-19], Debnath et al. [11], Velmurugan et al. [20]* and many others.

## II. THE TRIPLE PASCAL MATRIX OF INVERSE AND TRIPLE PASCAL SEQUENCE SPACES

Let  $P$  denote the Pascal means defined by the Pascal matrix as is defined by

$$\text{if } 0 \leq m \leq r, 0 \leq n \leq s, 0 \leq k \leq t$$

$$\text{if } m > r, n > s, k > t; r, s, t, m, n, k \in \mathbb{N}$$

and the inverse of Pascal's matrix

$$P = [P_{mnk}^{rst}]^{-1} = \begin{cases} (-1)^{(r-m)+(s-n)+(t-k)} \binom{r}{m} \binom{s}{n} \binom{t}{k} & \text{if } 0 \leq m \leq r, 0 \leq n \leq s, 0 \leq k \leq t \\ 0, \text{ if } (m > r, n > s, k > t; r, s, t, m, n, k \in \mathbb{N}) \end{cases}$$

... (\*)

There is some interesting properties of Pascal matrix. For example, we can form three types of matrix; symmetric, lower triangular and upper triangular; for any integer  $i, j, k > 0$ . The symmetric Pascal matrix of order  $n \times n \times n$  is defined by

$$A_{ijk} = a_{ijk} = \binom{i+j+k}{i} \binom{i+j+k}{j} \binom{i+j+k}{k} \text{ for } i, j, k = 0, 1, 2, \dots, n. \quad (5)$$

We can define the lower triangular Pascal matrix of order  $n \times n \times n$  by

$$L_{ijk} = (L_{ijk}) = \frac{1}{(-1)^{i+j-2k} I_{ijk}}; i, j, k = 1, 2, \dots, n. \quad (6)$$

and the upper triangular Pascal matrix of order  $n \times n \times n$  is defined by

$$U_{ijk} = (U_{ijk}) = \frac{1}{(-1)^{k-(i+j)} I_{ijk}}; i, j, k = 1, 2, \dots, n. \quad (7)$$

We know that  $U_{ijk} = (L_{ijk})^T$  for any positive integer  $i, j, k$ .

(i) Let  $A_{ijk}$  be the symmetric Pascal matrix of order  $n \times n \times n$  defined by (2.1),  $L_{ijk}$  be the lower triangular Pascal matrix of order  $n \times n \times n$  defined by (2.2), then  $A_{ijk} = L_{ijk} U_{ijk}$  and  $\det(A_{ijk}) = 1$ .

(ii) Let  $A$  and  $B$  be  $n \times n \times n$  matrices. We say that  $A$  is similar to  $B$  if there is an invertible  $n \times n \times n$  matrix  $P$  such that  $P^{-1}AP = B$ .

(iii) Let  $A_{ijk}$  be the symmetric Pascal matrix of order  $n \times n \times n$  defined by (2.1), then  $A_{ijk}$  is similar to its inverse  $A_{ijk}^{-1}$ .

(iv) Let  $L_{ijk}$  be the lower triangular Pascal matrix of order  $n \times n \times n$  defined by (2.2), then  $L_n^{-1} = L_{ijk}^{-1} = (-1)^{i+j-2k} I_{ijk}$ .

We wish to introduce the Pascal sequence spaces  $P_{\Lambda^3}$  and  $P_{\chi^3}$  as the set of all sequences such that  $P$ -transforms of them are in the spaces  $\Lambda^3$  and  $\chi^3$ , respectively, that is

$$\Lambda_P^3 = \eta_{mnk} = \left\{ x = (x_{mnk}) : \sup_{rst} \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} |x_{mnk}|^{\frac{1}{m+n+k}} < \infty \right\}, \text{ and } \chi_P^3 = \mu_{mnk} = \left\{ x = (x_{mnk}) : \lim_{rst \rightarrow \infty} \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} = 0 \right\}.$$

We may redefine the spaces  $\Lambda_P^3, \chi_P^3$  as follows:  $\Lambda_P^3 = P_{\Lambda^3}, \chi_P^3 = P_{\chi^3}$ .

If  $\lambda$  is a normed or paranormed sequence space; then matrix domain  $\lambda_P$  is called a Pascal triple sequence space. We define the triple sequence  $y = (y_{rst})$  which will be frequently used, as the  $P$ -transform of a triple sequence  $x = (x_{rst})$  i.e.,

$$y_{rst} = \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} x_{mnk}, (r, s, t \in \mathbb{N}). \quad (8)$$

Pascal sequence spaces  $P_{\Lambda^3}$  and  $P_{\chi^3}$  as the set of all sequences such that  $P$ -transforms of them are in the spaces  $\Lambda^3$  and  $\chi^3$ ,

respectively, that is

It can be shown easily that  $P_{\chi^3}$  are linear and metric space by the following metric:

$$d(x, y)_{P_{\chi^3}} = d(Px, Py) = \sup_{m,n,k} \left\{ ((m+n+k)! |x_{mnk} - y_{mnk}|)^{\frac{1}{m+n+k}} : m, n, k = 1, 2, 3, \dots \right\}.$$

Consider a triple sequence  $x = (x_{mnk})$ . The  $(m, n, k)^{th}$  section  $x^{[m, n, k]}$  of the sequence is defined by  $x^{[m, n, k]} = \sum_{i, j, q=0}^{m, n, k} x_{ijq} \mathfrak{S}_{ijq}$  for all  $m, n, k \in \mathbb{N}$ ; where  $\mathfrak{S}_{ijq}$  denotes the triple sequence whose only non zero term is a  $\frac{1}{(i+j+k)!}$  in the  $(i, j, k)^{th}$  place for each  $i, j, k \in \mathbb{N}$ .

If  $X$  is a sequence space, we give the following definitions:

(i)  $X'$  is continuous dual of  $X$ ;

$$(ii) X^\alpha = \left\{ a = (a_{mnk}) : \sum_{m,n,k=1}^{\infty} |a_{mnk} x_{mnk}| < \infty, \text{ for each } x \in X \right\};$$

$$(iii) X^\beta = \left\{ a = (a_{mnk}) : \sum_{m,n,k=1}^{\infty} a_{mnk} x_{mnk} \text{ is convergent, for each } x \in X \right\};$$

$$(iv) X^\gamma = \left\{ a = (a_{mnk}) : \sup_{m,n,k \geq 1} \left| \sum_{m,n,k=1}^{M,N,K} a_{mnk} x_{mnk} \right| < \infty, \text{ for each } x \in X \right\};$$

$$(v) \text{ Let } X \text{ be an FK-space } \supset \phi; \text{ then } X^f = \left\{ f(\mathfrak{S}_{mnk}) : f \in X' \right\};$$

$$(vi) X^\delta = \left\{ a = (a_{mnk}) : \sup_{m,n,k} |a_{mnk} x_{mnk}|^{1/m+n+k} < \infty, \text{ for each } x \in X \right\};$$

$X^\alpha, X^\beta, X^\gamma$  are called  $\alpha$  - (or Köthe - Toeplitz) dual of  $X$ ,  $\beta$  - (or generalized - Köthe - Toeplitz) dual of  $X$ ,  $\gamma$  - dual of  $X$ ,  $\delta$  - dual of  $X$  respectively.  $X^\alpha$  is defined by Gupta and Kamptan [21]. It is clear that  $X^\alpha \subset X^\beta$  and  $X^\alpha \subset X^\gamma$ , but  $X^\alpha \subset X^\gamma$  does not hold.

### III. MAIN RESULTS

#### A. Theorem

The triple sequence spaces  $P_{\chi^3}$  and  $P_{\Lambda^3}$  are linearly isomorphic spaces  $\Lambda^3$  and  $\chi^3$  respectively i.e.,  $P_{\Lambda^3} \cong \Lambda^3$  and  $P_{\chi^3} \cong \chi^3$ .

Proof: To prove the fact  $P_{\chi^3} \cong \chi^3$ , we should show the existence of a linear bijection between the spaces  $P_{\chi^3}$  and  $\chi^3$ . Consider the transformation  $T$  defined with the notation (2.4), from  $P_{\chi^3}$  to  $\chi^3$ . The linearity to  $T$  is clear. Further, it is trivial that  $T\mu = 0$  whenever  $T\mu = 0$  and hence  $T$  is injective.

Let  $\gamma \in \chi^3$ . We consider the triple sequence  $\mu = (\mu_{mnk})$  as follows:

$$\mu_{mnk} = \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t (-1)^{(r-m)+(s-n)+(t-k)} \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |y_{mnk}|)^{\frac{1}{m+n+k}}.$$

Then

$$\begin{aligned} \lim_{r,s,t \rightarrow \infty} (Px)_{rst} &= \lim_{r,s,t \rightarrow \infty} \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \\ &(-1)^{(r-m)+(s-n)+(t-k)} \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |y_{mnk}|)^{\frac{1}{m+n+k}} = 0. \end{aligned}$$

Thus, we have that  $x \in P_{\chi^3}$ . In addition note that

$$\begin{aligned} d(x, y)_{P_{\chi^3}} &= \sup_{r,s,t} \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \\ &(-1)^{(r-m)+(s-n)+(t-k)} \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk} - y_{mnk}|)^{\frac{1}{m+n+k}} \end{aligned}$$

$$\begin{aligned}
&= \sup_{r,s,t} ((r+s+t)! |x_{rst} - y_{rst}|)^{\frac{1}{r+s+t}} \\
&= d(x,y)_{\chi^3} < \infty.
\end{aligned}$$

Consequently,  $T$  is surjective and is metric preserving. Hence,  $T$  is a linear bijection which therefore says us that the spaces  $P_{\chi^3}$  to  $\chi^3$  are linearly isomorphic. In the same way, it can be shown that  $P_{\Lambda^3}$  are linearly isomorphic to  $\Lambda^3$ , respectively and so we omit the detail.

### B. Theorem

Let  $(m, n, k) \in \mathbb{N}^3$  be a fixed triple number and  $b^{(mnk)} = \left\{ b_{rst}^{(mnk)} \right\}_{(r,s,t) \in \mathbb{N}^3}$ , where

$$b_{rst}^{(mnk)} = \begin{cases} 0, & \text{if } 0 \leq r \leq m, s \leq n, t \leq k \\ (-1)^{(r-m)+(s-n)+(t-k)} \binom{r}{m} \binom{s}{n} \binom{t}{k} & \text{if } r \geq m, s \geq n, t \geq k. \end{cases}$$

Then the following assertions are true:

(i) The triple sequence  $\left\{ b_{rst}^{(mnk)} \right\}$  is a basis for the space  $P_{\chi^3}$  and every  $x \in P_{\chi^3}$  has a unique representation of the form  $\mu = \sum_m \sum_n \sum_k \lambda_{mnk} b^{(mnk)}$ , where  $\lambda_{mnk} = \left( P((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} \right)_{m,n,k}$  for all  $m, n, k \in \mathbb{N}$ .

### C. Proposition

The triple sequence  $P_{\chi^3}$  is a linear set over the set of complex numbers  $\mathbb{C}$ .

**Proof:** It is trivial. Therefore, the proof is omitted.

### D. Proposition

$$\left( P_{\chi^3} \right)^\delta \not\subset P_{\Lambda^3}$$

**Proof:** Let  $\gamma \in \delta$ - dual of  $P_{\chi^3}$ . Then  $|\mu_{mnk} \gamma_{mnk}| \leq M^{m+n+k}$  for some constant  $M > 0$  and for each  $\mu \in P_{\chi^3}$ . Therefore,  $|\gamma_{mnk}| \leq M^{m+n+k}$  for each  $m, n, k$  by taking  $\mu = (\mathfrak{S}_{mnk})$ . This implies that  $\gamma \in P_{\Lambda^3}$ . Thus,

$$\left( P_{\chi^3} \right)^\delta \subset P_{\Lambda^3} \tag{9}$$

we now choose the triple sequences  $(\gamma_{mnk})$  and  $(\mu_{mnk})$  by  $(\gamma_{mnk}) = 1$  for all  $m, n$  and  $k$ , and by

$$\begin{aligned}
(m+2)! x_{m11} &= \left[ 2 \frac{r!}{m!(r-m)!} \frac{s!}{1!(s-1)!} \frac{t!}{1!(t-k)!} \right]^{(m+2)^2} \text{ and} \\
(m+n+k)! x_{mnk} &= \left[ 2 \frac{r(r-1)!}{m(m-1)!(r-m)!} \frac{s(s-1)!}{n(n-1)!(s-n)!} \frac{t(t-1)!}{k(k-1)!(t-k)!} \right] = 0(r, s, t = 0) \text{ for all } m, n, k = 0
\end{aligned}$$

Obviously,  $\gamma \in P_{\Lambda^3}$  and since  $(m+n+k)! x_{mnk} = 0$  for all  $m, n, k = 0$ ,

$(m+n+k)! x_{mnk}$  converges to zero. Hence,  $\mu \in P_{\chi^3}$ . But

$$((m+2)! |a_{m11} x_{m11}|)^{\frac{1}{m+n+k}} = \left[ 2 \frac{r!}{m!(r-m)!} \frac{s!}{1!(s-1)!} \frac{t!}{1!(t-1)!} \right]^{(m+2)^2} \rightarrow \infty \text{ as } m \rightarrow \infty, \text{ hence}$$

$$\mu \notin (P_{\chi^3})^\delta \quad (10)$$

From (3.1) and (3.2), we are granted  $(P_{\chi^3})^\delta \not\subseteq P_{\chi^3}$ . This completes the proof.

### E. Proposition:

The dual space of  $P_{\chi^3}$  is  $P_{\chi^3}$ . In other words  $(P_{\chi^3})^* = P_{\chi^3}$ .

**Proof:** We recall that  $\mathfrak{S}_{mnk} =$

$$\begin{pmatrix} 0 & 0 & \dots & 0 & \dots \\ 0 & 0 & \dots & 0 & \dots \\ \cdot & & & & \\ \cdot & & & & \\ \cdot & & & & \\ 0 & 0 & \dots & \frac{1}{(m+n+k)!} & 0 & \dots \\ 0 & 0 & \dots & 0 & 0 & \dots \end{pmatrix}$$

with  $\frac{1}{(m+n+k)!}$  in the  $(m, n, k)$ th position and zero's else where, with  $\mu = \mathfrak{S}_{mnk}$ ,

which is a triple Pascal chi sequence. Hence,  $\mathfrak{S}_{mnk} \in P_{\chi^3}$ ;  $f(x) = \sum_{m,n,k=1}^{\infty} \mu_{mnk} \gamma_{mnk}$  with  $\mu \in P_{\chi^3}$  and  $f \in (P_{\chi^3})^*$ , where  $(P_{\chi^3})^*$  is the dual space of  $P_{\chi^3}$ . Take  $\mu = (\mu_{mnk}) = \mathfrak{S}_{mnk} \in P_{\chi^3}$ . Then,

$$|\gamma_{mnk}| \leq \|f\| d(\mathfrak{S}_{mnk}, 0) < \infty \quad \forall m, n, k. \quad (11)$$

Thus,  $(\gamma_{mnk})$  is a bounded sequence and hence an triple Pascal analytic sequence. In other words,  $\gamma \in P_{\Lambda^3}$ . Therefore  $(P_{\chi^3})^* = P_{\Lambda^3}$ . This completes the proof.

### F. Proposition:

$$(P_{\Lambda^3})^\beta \not\subseteq P_{\chi^3}$$

**Proof: Step 1:** Let  $(\mu_{mnk}) \in (P_{\Lambda^3})^\beta$ ,

$$\sum_{m,n,k=1}^{\infty} |\mu_{mnk} \gamma_{mnk}| < \infty \quad \forall (\gamma_{mnk}) \in P_{\Lambda^3} \quad (12)$$

Let us assume that  $(\mu_{mnk}) \notin P_{\chi^3}$ . Then, there exists a strictly increasing sequence of positive integers  $(m_p + n_p + k_p)$  such that

$$(m_p + n_p + k_p)! \left| x_{(m_p + n_p + k_p)} \right| > \frac{1}{\left[ 2 \frac{r!}{m!(r-m)!} \frac{s!}{1!(s-1)!} \frac{t!}{1!(t-k)!} \right] (m_p + n_p + k_p)}, \quad (p = 1, 2, 3, \dots) \quad (13)$$

Let

$$(m_p + n_p + k_p)! \gamma_{(m_p + n_p + k_p)} = \left[ 2 \frac{r!}{m!(r-m)!} \frac{s!}{1!(s-1)!} \frac{t!}{1!(t-k)!} \right]^{(m_p + n_p + k_p)} \text{ for } (p = 1, 2, 3, \dots)$$

$$\gamma_{mnk} = 0 \text{ otherwise}$$

Then,  $(\gamma_{mnk}) \in P_{\Lambda^3}$ . However,

$$\sum_{m,n,k=1}^{\infty} |\mu_{mnk} \gamma_{mnk}| = \sum_{p=1}^{\infty} (m_p + n_p + k_p)! \left| \mu_{(m_p n_p k_p)} \gamma_{(m_p n_p k_p)} \right| > 1 + 1 + \dots$$

We know that the infinite series  $1+1+1+\dots$  diverges. Hence  $\sum_{m,n,k=1}^{\infty} |\mu_{mnk} \gamma_{mnk}|$  diverges. This contradicts (3.4). Hence  $(\mu_{mnk}) \in P_{\chi^3}$ . Therefore,

$$(P_{\Lambda^3})^\beta \subset P_{\chi^3} \quad (14)$$

and  $\gamma_{mnk} = \mu_{mnk} = 0 (m > 1)$  for all  $n, k$  then obviously  $\mu \in P_{\chi^3}$  and  $\gamma \in P_{\Lambda^3}$ , but  $\sum_{m,n,k=1}^{\infty} \mu_{mnk} \gamma_{mnk} = \infty$ . Hence,

$$\gamma \notin (P_{\Lambda^3})^\beta \quad (15)$$

From (3.6) and (3.7), we are granted  $(P_{\Lambda^3})^\beta \not\subset P_{\chi^3}$ . This completes the proof.

### G. Definition

Let  $p = (p_{mnk})$  be a triple sequence of positive real numbers. Then,

$$P_{\chi^3}(p) = \left( \lim_{r,s,t \rightarrow \infty} \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} = 0 \right)^{p_{mnk}}. \quad (16)$$

Suppose that  $p_{mnk}$  is a constant for all  $m, n, k$  then  $P_{\chi^3}(p) = P_{\chi^3}$ .

### H. Proposition

Let  $0 \leq p_{mnk} \leq q_{mnk}$  for all  $m, n, k \in \mathbb{N}$  and let  $\left\{ \frac{q_{mnk}}{p_{mnk}} \right\}$  be bounded. Then  $P_{\chi^3}(q) \subset P_{\chi^3}(p)$ .

**Proof:** Let

$$\mu \in P_{\chi^3}(q), \text{ then} \quad (17)$$

$$\left( \lim_{r,s,t \rightarrow \infty} \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} = 0 \right)^{q_{mnk}}. \quad (18)$$

Let  $t_{mnk} = \left( \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} \right)^{q_{mnk}}$ , and let  $\alpha_{mnk} = p_{mnk}/q_{mnk}$ . Since  $p_{mnk} \leq q_{mnk}$ , we have  $0 \leq \alpha_{mnk} \leq 1$ . Let  $0 < \alpha < \alpha_{mnk}$ . then

$$u_{mnk} = \begin{cases} t_{mnk}, & \text{if } (t_{mnk} \geq 1) \\ 0, & \text{if } (t_{mnk} < 1) \end{cases}$$

$$v_{mnk} = \begin{cases} 0, & \text{if } (t_{mnk} \geq 1) \\ t_{mnk}, & \text{if } (t_{mnk} < 1) \end{cases} \quad (19)$$

$$t_{mnk} = u_{mnk} + v_{mnk}, t_{mnk}^{\alpha_{mnk}} = u_{mnk}^{\alpha_{mnk}} + v_{mnk}^{\alpha_{mnk}}.$$

Now, it follows that

$$u_{mnk}^{\alpha_{mnk}} \leq u_{mnk} \leq t_{mnk}, v_{mnk}^{\alpha_{mnk}} \leq u_{mnk}^{\alpha_{mnk}} \quad (20)$$

Since  $t_{mnk}^{\alpha_{mnk}} = u_{mnk}^{\alpha_{mnk}} + v_{mnk}^{\alpha_{mnk}}$ , we have  $t_{mnk}^{\alpha_{mnk}} \leq t_{mnk} + v_{mnk}^{\alpha_{mnk}}$ . Thus,

$$\begin{aligned} & \left( \left( \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} \right)^{q_{mnk}} \right)^{\alpha_{mnk}} \leq \\ & \left( \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} \right)^{q_{mnk}} \\ & \left( \left( \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} \right)^{q_{mnk}} \right)^{(p_{mnk}/q_{mnk})} \leq \\ & \left( \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} \right)^{q_{mnk}} \dots * \end{aligned}$$

which yields

$$\begin{aligned} & \left( \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} \right)^{p_{mnk}} \\ & \leq \left( \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} \right)^{q_{mnk}}. \end{aligned}$$

However,  $\left( \lim_{rst \rightarrow \infty} \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} = 0 \right)^{q_{mnk}}$  (by(3.10)). Thus,

$$\left( \lim_{rst \rightarrow \infty} \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} = 0 \right)^{p_{mnk}}.$$

Hence,

$$\mu \in P_{\mathcal{X}^3}(p). \quad (21)$$

Hence (3.9) and (3.13), we are granted

$$P_{\mathcal{X}^3}(q) \subset P_{\mathcal{X}^3}(p). \quad (22)$$

This completes the proof.



**I. Proposition**

(a) Let  $0 < \inf p_{mnk} \leq p_{mnk} \leq 1$ , then  $P_{\chi^3}(p) \subset P_{\chi^3}$ .

(b) If  $1 \leq p_{mnk} \leq \sup p_{mnk} < \infty$ , then  $P_{\chi^3} \subset P_{\chi^3}(p)$ .

**Proof:** The above statements are special cases of proposition 3.8. Therefore, it can be proved by similar arguments.

**J. Proposition**

If  $0 < p_{mnk} \leq q_{mnk} < \infty$  for each  $m, n, k$  then  $P_{\chi^3}(p) \subseteq P_{\chi^3}(q)$ .

**Proof:** Let  $\mu \in P_{\chi^3}(p)$ , then

$$\left( \lim_{r,s,t \rightarrow \infty} \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} = 0 \right)^{p_{mnk}} \tag{23}$$

$$\left( \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} \right)^{q_{mnk}} \leq$$

$$\left( \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} \right)^{p_{mnk}} \dots \dots *, \text{ then}$$

$$\left( \lim_{rst \rightarrow \infty} \sum_{m=0}^r \sum_{n=0}^s \sum_{k=0}^t \binom{r}{m} \binom{s}{n} \binom{t}{k} ((m+n+k)! |x_{mnk}|)^{\frac{1}{m+n+k}} = 0 \right)^{q_{mnk}} \text{ (by using (3.15)). We have } \mu \in P_{\chi^3}(q).$$

Hence,  $P_{\chi^3}(p) \subseteq P_{\chi^3}(q)$ . This completes the proof.

**Competing Interests:** The authors declare that there is not any conflict of interests regarding the publication of this manuscript.

[1] Bipan Hazarika, N. Subramanian and A. Esi, On rough weighted ideal convergence of triple sequence of Bernstein polynomials, *Proceedings of the Jangjeon Mathematical Society*, **21(3)** (2018), 497-506.

[2] A. Esi, On some triple almost lacunary sequence spaces defined by Orlicz functions, *Research and Reviews: Discrete Mathematical Structures*, **1(2)**, (2014), 16-25.

[3] A. Esi and M. Necdet Catalbas, Almost convergence of triple sequences, *Global Journal of Mathematical Analysis*, **2(1)**, (2014), 6-10.

[4] A. Esi and E. Savas, On lacunary statistically convergent triple sequences in probabilistic normed space, *Appl. Math. and Inf. Sci.*, **9(5)**, (2015), 2529-2534.

[5] A. Esi, S. Araci and M. Acikgoz, Statistical Convergence of Bernstein Operators, *Appl. Math. and Inf. Sci.*, **10(6)**, (2016), 2083-2086.

[6] A. Esi, S. Araci and Ayten Esi,  $\lambda$ -Statistical Convergence of Bernstein polynomial sequences, *Advances and Applications in Mathematical Sciences*, **16(3)**, (2017), 113-119.

[7] A. Esi, N. Subramanian and Ayten Esi, On triple sequence space of Bernstein operator of rough  $I$ -convergence Pre-Cauchy sequences, *Proyecciones Journal of Mathematics*, **36(4)**, (2017), 567-587.

[8] A. Esi and N. Subramanian, Generalized rough Cesaro and lacunary statistical Triple difference sequence spaces in probability of fractional order defined by Musielak Orlicz function, *International Journal of Analysis and Applications*, **16(1)** (2018), 16-24.

[9] A. Esi and N. Subramanian, On triple sequence spaces of Bernstein operator of  $\chi^3$  of rough  $\lambda$ -statistical convergence in probability of random variables defined by Musielak-Orlicz function, *Int. J. open problems Compt. Math*, **11(2)** (2019), 62-70.

[10] A. J. Dutta A. Esi and B.C. Tripathy, Statistically convergent

- triple sequence spaces defined by Orlicz function , *Journal of Mathematical Analysis*, **4(2)**, (2013), 16-22.
- [11] S. Debnath, B. Sarma and B.C. Das ,Some generalized triple sequence spaces of real numbers, *Journal of nonlinear analysis and optimization*, **Vol. 6, No. 1** (2015), 71-79.
- [12] A. Sahiner, M. Gurdal and F.K. Duden, Triple sequences and their statistical convergence, *Selcuk J. Appl. Math.* , **8 No. (2)**(2007), 49-55.
- [13] A. Sahiner, B.C. Tripathy , Some  $I$  related properties of triple sequences, *Selcuk J. Appl. Math.*, **9 No. (2)**(2008), 9-18.
- [14] N. Subramanian and A. Esi, The generalized triple difference of  $\chi^3$  sequence spaces, *Global Journal of Mathematical Analysis*, **3 (2)** (2015), 54-60.
- [15] N. Subramanian and A. Esi, Rough Variables of convergence, *Vasile Alecsandri University of Bacau Faculty of Sciences, Scientific studies and Research series Mathematics and informatics*, **27 (2)** (2017), 65-72.
- [16] N. Subramanian and A. Esi, Wijsman rough convergence triple sequences, *Matematychni studii*, **48 (2)** (2017), 171-179.
- [17] N. Subramanian and A. Esi, On triple sequence space of Bernstein operator of  $\chi^3$  of rough  $\lambda$ - statistical convergence in probability defined by Musielak-Orlicz function  $p$ - metric, *Electronic Journal of Mathematical Analysis and Applications*, **6 (1)** (2018), 198-203.
- [18] N. Subramanian, A. Esi and M. Kemal Ozdemir, Rough Statistical Convergence on Triple Sequence of Bernstein Operator of Random Variables in Probability , *Songklanakarin Journal of Science and Technology*, **in press** (2018).
- [19] N. Subramanian, A. Esi and V.A. Khan, The Rough Intuitionistic Fuzzy Zweier Lacunary Ideal Convergence of Triple Sequence spaces, *journal of mathematics and statistics*, **14** (2018), 72-78.
- [20] S. Velmurugan and N. Subramanian, Bernstein operator of rough  $\lambda$ - statistically and  $\rho$  Cauchy sequences convergence on triple sequence spaces, *Journal of Indian Mathematical Society*, **85 (1-2)** (2018), 257-265.
- [21] P.K.Kamthan and M.Gupta, Sequence spaces and series, Lecture notes, Pure and Applied Mathematics, *65 Marcel Dekker, Inc., New York* , 1981.